

Abstract Submitted
for the MAR13 Meeting of
The American Physical Society

Topological

Excitonic Superfluids in Three Dimensions¹ MATTHEW GILBERT, University of Illinois - Urbana-Champaign, EWELINA HANKIEWICZ, Universitaet Wuerzburg, YOUNGSEOK KIM, University of Illinois - Urbana-Champaign — We study the equilibrium and non-equilibrium properties of topological dipolar intersurface exciton condensates within time-reversal invariant topological insulators in three spatial dimensions without a magnetic field. We elucidate that, in order to correctly identify the proper pairing symmetry within the condensate order parameter, the full three-dimensional Hamiltonian must be considered. As a corollary, we demonstrate that only particles with similar chirality play a significant role in condensate formation. Furthermore, we find that the intersurface exciton condensation is not suppressed by the interconnection of surfaces in three-dimensional topological insulators as the intersurface polarizability vanishes in the condensed phase. This eliminates the surface current flow leaving only intersurface current flow through the bulk. We conclude by illustrating how the excitonic superfluidity may be identified through an examination of the terminal currents above and below the condensate critical current.

¹Army Research Office (ARO) under contract number W911NF-09-1-0347, the Office of Naval Research (ONR) under contract number N0014-11-1-0728, and the Air Force Office of Scientific Research (AFOSR) under contract number FA9550-10-1-0459, DFG Grant HA 5893

Matthew Gilbert
University of Illinois - Urbana-Champaign

Date submitted: 07 Nov 2012

Electronic form version 1.4