

Abstract Submitted
for the MAR13 Meeting of
The American Physical Society

What is the granular response to a high-speed impact?¹ ABRAM CLARK, Duke University, LOU KONDIC, New Jersey Institute of Technology, ROBERT BEHRINGER, Duke University — Although many studies of impact on a granular material exist, the connections between the local granular response, the microscopic processes which dissipate kinetic energy, and the intruder dynamics are unclear, largely due to experimental difficulties in obtaining very fast data at the grain scale. We use high-speed imaging (40 kHz) of an intruder striking a quasi-2D system of photoelastic disks, yielding both the intruder dynamics and the force response of individual grains. The frame rates are fast enough to resolve rich acoustic activity on the particle scale. For long time scales, the intruder dynamics are consistent with previously used empirical force laws. However, for short time scales, we observe very large fluctuations in the deceleration, which we connect to the intermittent acoustic activity beneath the intruder as it moves. We show that these intense, intermittent acoustic pulses, which travel much faster than the intruder along networks of grains, are the primary microscopic mechanism of energy loss. These pulses carry energy away into the medium, and they decay roughly exponentially with distance. We examine the statistics of these fluctuations in order to better understand their origin and behavior.

¹Supported by the US DTRA under grant HDTRA1-10-0021

Abram Clark
Duke University

Date submitted: 07 Nov 2012

Electronic form version 1.4