

Abstract Submitted
for the MAR13 Meeting of
The American Physical Society

Very Narrow Intersubband Excitations in High Mobility 2DESs¹

URSULA WURSTBAUER, ARON PINCZUK, Columbia University, JOHN WATSON, SUMIT MONDAL, MICHAEL J. MANFRA, Purdue University, KEN W. WEST, LOREN N. PFEIFFER, Princeton University — We report the observation of very narrow collective intersubband excitations (ISBE) of 2D electron systems (2DESs) with ultra-high mobility ($\mu \geq 15 \times 10^6$ cm²/Vs) in high quality GaAs quantum structures. These findings from resonant inelastic light scattering (RILS) experiments are used as tools for exploration of links between transport mobility and collective electron behavior in 2DES of high perfection. We find that the line-widths of collective ISB modes can be as low as $80 \mu\text{eV}$. Comparison of ISBE measurements from several samples exhibits a variation in line-width of more than a factor of two. There is, however, a surprising lack of direct correlation between ISBE line-width with mobility in the range $15 \geq \mu \geq 24 \times 10^6$ cm²/Vs. Measurements of ISBE by RILS will be evaluated as a method to explore the interplay of quality (as indicated by mobility) and fundamental interactions in the fractional quantum Hall effect.

¹Supported by NSF and AvH

Ursula Wurstbauer
Columbia University

Date submitted: 09 Nov 2012

Electronic form version 1.4