

Abstract Submitted  
for the MAR13 Meeting of  
The American Physical Society

**Exciton-plasmon coupling in monolayer molybdenum disulfide<sup>1</sup>**

JED ZIEGLER, A.K.M. NEWAZ, KIRILL BOLOTIN, RICHARD HAGLUND, Vanderbilt University — Two-dimensional materials such as monolayer molybdenum disulfide ( $\text{MoS}_2$ ) represent a unique platform for investigating the dynamics of exciton-plasmon coupling. We report on the generation and modulation of coherent and incoherent coupled states between excitons in monolayer  $\text{MoS}_2$  and plasmons in an array of gold nanoparticle deposited onto the surface of  $\text{MoS}_2$ . We study the behavior of these coherent states, termed plexcitons using a combination of photoluminescence, extinction and ultrafast spectroscopies. The close proximity of the two characteristic exciton bands of  $\text{MoS}_2$  presents multiple coherent coupling configurations, including A-or-B exciton-plasmon, and A-and-B exciton-plasmon interactions. These configurations of plexciton formation that are shown to modulate both the extinction and photoluminescence spectra of the hybrid system. This includes broadband photoluminescence and Fano-type resonances. This behavior is distinct from the spectral response of the  $\text{MoS}_2$  and plasmonic components of the system. Incoherent exciton-plasmon coupling, achieved by detuning from the plasmon extinction peaks, enhances the interaction of  $\text{MoS}_2$  with light by focusing the plasmon energy. Depending on which coupling configuration is chosen, our results show that the  $\text{MoS}_2$ /plasmon hybrid systems can act as high efficiency light harvesters, broadband emitters and as tunable visible and NIR photodetectors.

<sup>1</sup>Support by Defense Threat Reduction Agency (HDTRA1-1-10-1-0047) and NSF DMR-1056859

Jed Ziegler  
Vanderbilt University

Date submitted: 18 Dec 2012

Electronic form version 1.4