

Abstract Submitted
for the MAR14 Meeting of
The American Physical Society

New Phase Transition in High Pressure Molecular Oxygen

YANIER CRESPO, SANDRO SCANDOLO, The Abdus Salam ICTP, Strada Costiera 11, I-34151 Trieste, Italy, ERIO TOSATTI, SISSA, Via Bonomea 265, I-34136 Trieste, Italy — Oxygen, one of the most common and important elements in nature, has an exceedingly well explored phase diagram under pressure, up and beyond 100 GPa. We propose a subtle, yet surprisingly undetected pressure-induced phase transition of molecular O_2 within the ϵ phase, near 20 GPa. While thus far in the whole ϵ phase region from about 8 to 96 GPa each individual O_2 molecule was generally believed to be in an $S=0$ state, we provide theoretical results, strongly connecting with existing vibrational and structural evidence, showing that this holds only above 20 GPa, whereas there is a collective switch from $S=0$ to an $S=1$ molecular state at and below 20 GPa. The former ϵ phase thus breaks up into two: a spinless ϵ_0 (20-96 GPa), and ϵ_1 (8-20 GPa) where the molecules possess spin with at least short range antiferromagnetic correlations.

Erio Tosatti
SISSA, Via Bonomea 265, I-34136 Trieste, Italy

Date submitted: 31 Oct 2013

Electronic form version 1.4