

Abstract Submitted
for the MAR14 Meeting of
The American Physical Society

Determination of Surface-Substrate Adsorption Energy using the Exchange-Hole Dipole Moment MATTHEW CHRISTIAN, ALBERTO OTERO DE LA ROZA, ERIN JOHNSON, Univ of California - Merced — Calculated surface-substrate binding energies are usually underestimated because conventional density functionals do not include dispersion, which is necessary to capture the van der Waals interactions that lead to weak physisorption. The exchange-hole dipole moment (XDM) model is a non-empirical density-functional approach to model dispersion. Adsorption energies for several aromatic molecules and nucleobases on noble metal surfaces were calculated using B86bPBE-XDM. In this talk, I compare the calculated adsorption energies with experiment and present implications for future applications to modeling surface interactions.

- [1] A. Otero-de-la-Roza and E. R. Johnson, *J. Chem. Phys.* **138** 204109 (2013).
- [2] A. Otero-de-la-Roza and E. R. Johnson, *J. Chem. Phys.* **137** 054103 (2012).
- [3] A. Otero-de-la-Roza and E. R. Johnson, *J. Chem. Phys.* **136** 204109 (2012).

Matthew Christian
Univ of California - Merced

Date submitted: 11 Nov 2013

Electronic form version 1.4