

Abstract Submitted
for the MAR14 Meeting of
The American Physical Society

Liquid drops on soft solids LUUK A. LUBBERS, JOOST H. WEIJS, University of Twente, SIDDHARTHA DAS, University of Alberta, LORENZO BOTTO, Imperial College, BRUNO ANDREOTTI, University Paris-Diderot, JACCO H. SNOEIJER, University of Twente — A sessile drop can elastically deform a substrate by the action of capillary forces. The typical size of the deformation is given by the ratio of surface tension and the elastic modulus, γ/E , which can reach up to 10-100 microns for soft elastomers. In this talk we theoretically show that the contact angles of drops on such a surface exhibit two transitions when increasing γ/E : (i) the microscopic geometry of the contact line first develops a Neumann-like cusp when γ/E is of the order of few nanometers, (ii) the macroscopic angle of the drop is altered only when γ/E reaches the size of the drop. Using the same framework we then show that two neighboring drops exhibit an effective interaction, mediated by the deformation of the elastic medium. This is in analogy to the well-known Cheerios effect, where small particles at a liquid interface attract each other due to the meniscus deformations. Here we reveal the nature of drop-drop interactions on a soft substrate by combining numerical and analytical calculations.

Luuk Lubbers
University of Twente

Date submitted: 14 Nov 2013

Electronic form version 1.4