

Abstract Submitted
for the MAR14 Meeting of
The American Physical Society

Self-assembly of Model Microtubules: Shape, Chirality and Twist

MARK STEVENS, Sandia National Labs, SHENGFENG CHENG, Virginia Polytechnic and State University — The efficient and controlled assembly of complex structures from macromolecular building blocks is a critical open question. Microtubules are one example of a biopolymer that possesses characteristics quite distinct from standard synthetic polymers that are derived from its monomer being a protein. In order to understand microtubule assembly and how to design artificial polymers that possess features similar to those of microtubules, we have used molecular dynamics simulations to study the self-assembly of model monomers into a tubule geometry. The self-assembly of free monomers into tubules yields a tubule pitch that often does not match the chirality of the monomer (including achiral monomers). We show that this mismatch occurs because of a twist deformation that brings the lateral interaction sites into alignment when the tubule pitch differs from the monomer chirality. In order to control the tubule pitch by preventing the twist deformation, we employ a lock-and-key interaction and obtain good control of the self-assembled tubule pitch. These results explain some fundamental features of microtubules. We generally find that the control of the assembly is limited, which explains the range of pitch and protofilament number observed in microtubule assembly.

Mark Stevens
Sandia National Labs

Date submitted: 14 Nov 2013

Electronic form version 1.4