

Abstract Submitted
for the MAR14 Meeting of
The American Physical Society

Finite-temperature effects in rotational hysteresis of ring Bose-Einstein condensates¹ N. MURRAY, C. LANIER, M. EDWARDS, Georgia Southern University, Y.-H. WANG, C.W. CLARK, S. ECKEL, F. JENDRZEJSKI, G.K. CAMPBELL, Joint Quantum Institute — A ring Bose-Einstein condensate (BEC) with zero circulation ($m = 0$) stirred by a barrier will eventually jump to an $m = 1$ state when stirred faster than a certain critical speed, Ω_c^+ . A ring BEC with $m = 1$ will drop to $m = 0$ when stirred at a critical speed, Ω_c^- , which is lower than Ω_c^+ . The loop areas, $\Omega_c^+ - \Omega_c^-$, of this hysteretic response of the BEC to stirring predicted by zero-temperature Gross-Pitaevskii equation (GPE) disagreed significantly with the results of a recent experiment. In the work reported here, we simulated this experiment with the phenomenologically damped GPE, [S. Choi, S. A. Morgan, and K. Burnett, Phys. Rev. A **57**, 4057 (1999)], and with the Zaremba-Nikuni-Griffin (ZNG) theory. The ZNG theory can account for finite-T, non-equilibrium dynamics. We compare the results of these simulations with the experimental data. The simulations show that a vortex/antivortex pair forms in the barrier region during the stirring and that this drives the hysteresis. We also show how the presence of an interacting, thermal cloud affects the dynamics of these pairs.

¹Supported in part by NSF grant #1068761 and ARO Atomtronics MURI

Mark Edwards
Georgia Southern Univ

Date submitted: 14 Nov 2013

Electronic form version 1.4