

Abstract Submitted  
for the MAR14 Meeting of  
The American Physical Society

**THz investigations of graphene-complex-oxide heterostructures<sup>1</sup>**

GIRIRAJ JNAWALI, LU CHEN, PATRICK IRVIN, JEREMY LEVY, University of Pittsburgh, SANGWOO RYU, CHANG-BEOM EOM, University of Wisconsin-Madison, FERESHTE GHAHARI, JAYAKANTH RAVICHANDRAN, PHILIP KIM, Columbia University — The unique and multifaceted properties of graphene have fascinated scientists and engineers for a decade now. A new frontier in research concerns properties of graphene in the THz-IR region, where the collective excitation of graphene 2D electron gas (2DEG) into plasmonic waves has proven the salient feature.<sup>2</sup> Complex oxide heterostructures (e.g., LaAlO<sub>3</sub>/SrTiO<sub>3</sub>, LAO/STO) also support a 2DEG with high carrier densities and expected plasmonic behavior. A unique feature of the LAO/STO system is the ability to control the electron density with nanoscale precision.<sup>3</sup> In addition, a method for sourcing and detecting broadband THz emission from LAO/STO nanojunctions has been recently demonstrated.<sup>4</sup> Here we describe initial efforts to investigate the THz properties of graphene-complex oxide (GCO) heterostructures. We envision that the proposed graphene plasmonic devices in the GCO will help to lay the foundation for a host of powerful THz-IR technologies for signal processing, imaging, spectroscopy and chemical sensing.

<sup>1</sup>We gratefully acknowledge support for this work from ONR (N00014-13-1-0806) and AFOSR (FA9550-12-1-0268).

<sup>2</sup>L. Ju, *et al.*, Nat. Nanotechnol. **6**, 630 (2011)

<sup>3</sup>C. Cen, *et al.*, Nat. Mater. **7**, 298 (2008)

<sup>4</sup>Y. Ma, *et al.*, Nano Lett. **13**, 2284 (2013)

Giriraj Jnawali  
University of Pittsburgh

Date submitted: 15 Nov 2013

Electronic form version 1.4