

Abstract Submitted
for the MAR14 Meeting of
The American Physical Society

Optical Detection of the Electron Nematic Phase in $\text{Sr}_3\text{Ru}_2\text{O}_7$

COLIN HEIKES, D. MACNEILL, Cornell University, S. GHOSH, S.N.Bose National Centre for Basic Sciences, R. PERRY, J.F. MERCURE, St. Andrews University, E.A. KIM, Cornell University, A. MACKENZIE, St. Andrews University, D.C. RALPH, Cornell University — We report the implementation of a fiber-based optical microscope, capable of operating at temperatures below 100 mK and in magnetic fields in excess of 9 Tesla, with sub-micron spatial resolution. This microscope is integrated into the bore of a dilution refrigerator with an optical fiber coupling light to an external optical table. Bench-top optical elements allow for polarization analysis of the reflected light from a surface and thus the detection of magnetic or other polarization-sensitive properties of mater at low temperature and high fields. We are studying the proposed electron nematic phase of the n=2 Ruddlesden-Popper material $\text{Sr}_3\text{Ru}_2\text{O}_7$, which exhibits a low-temperature phase transition in the form of an in-plane conduction anisotropy. We report recent results from concurrent transport measurements and polarization analysis as well as polarization microscopy with sample temperatures below 150 mK and applied magnetic fields from 0 T to 9 T.

Colin Heikes
Cornell University

Date submitted: 15 Nov 2013

Electronic form version 1.4