

Abstract Submitted  
for the MAR14 Meeting of  
The American Physical Society

**Intrinsic anisotropy of critical current in c-tilted MgB<sub>2</sub> thin films**

ANATOLII POLYANSKII, DMYTRO ABRAIMOV, FUMITAKE KAMETANI, DAVID LARBALESTIER, ASC, FSU, ALEX GUREVICH, Old Dominion University, AKI YAMAMOTO, University of Tokyo, I. PALLECCHI, MARINA PUTTI, University of Genova, C. ZHUANG, T. TAN, X.X. XI, Temple University — The intense investigations of MgB<sub>2</sub> in recent years have revealed many unusual effects of the two-band superconductivity on the properties of this material. However, given the size and geometry of available single crystals, the critical current along the c-direction, mostly affected by the anisotropy of the two bands, has not been thoroughly investigated. To address the issue of the c-axis current transport, we succeeded to measure  $J_c$  along the c axis by growing epitaxial films with tilted c-axis, which offers a unique possibility of probing both the in-plane and the out-of-plane critical currents. We show that anisotropic band parameters determine the anisotropy of vortex pinning with respect to the current direction. This yields a temperature and field dependent anisotropy of the critical current density  $J_c$  related to the anisotropic penetration depths at low temperature and low field and to the ratio of  $\sigma$  band effective masses along the c and ab crystalline axes at temperatures close to  $T_c$  or at applied fields larger than the virtual upper critical field of the  $\pi$  band  $H \approx 0.2T$ . Our results suggest that the out-of-plane current transport could set the ultimate  $J_c$  limit in randomly oriented MgB<sub>2</sub> polycrystals.

Anatolii Polyanskii  
ASC, FSU

Date submitted: 15 Nov 2013

Electronic form version 1.4