

MAR14-2013-020497

Abstract for an Invited Paper
for the MAR14 Meeting of
the American Physical Society

Colossal Enhancement of Spin-Orbit Coupling in Hydrogenated Graphene

BARBAROS OEZYILMAZ, National University of Singapore

Graphene's extremely small intrinsic spin-orbit (SO) interaction makes the realization of many interesting phenomena such as topological states and the spin Hall Effect (SHE) practically impossible. Recently, it was predicted that the introduction of adatoms in graphene would enhance the SO interaction by the conversion of sp² to sp³ bonds [1]. However, introducing adatoms and yet keeping graphene metallic, i.e., without creating electronic (Anderson) localization⁸ is experimentally challenging. Here, we show that the controlled addition of small amounts of covalently bonded hydrogen atoms is sufficient to induce a colossal enhancement of the SO interaction by three orders of magnitude. This results in a SHE at zero external magnetic field even at room temperature, with non-local spin signals up to 100 Ω . The SHE is, further, directly confirmed by the Larmor spin-precession measurements. From this and the length dependence of the non-local signal we extract a spin relaxation length $\sim 1 \mu\text{m}$, a spin relaxation time $\sim 90 \text{ ps}$ and a SO strength of 2.5 meV [2].

[1] Castro Neto, A. H. & Guinea, F. Impurity-induced spin-orbit coupling in graphene. *Phys. Rev. Lett.* (2009).

[2] Balakrishnan, J., Koon, G. K. W., Jaiswal, M., Castro Neto, A. H., Özyilmaz, B.; Colossal Enhancement of Spin-Orbit Coupling in Weakly Hydrogenated Graphene; *Nature Physics* (2013).