

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Phase diagram of a semimetal in the magnetic ultra-quantum limit¹ ALDO ISIDORI, FRANK ARNOLD, Royal Holloway, University of London, ERIK KAMPERT, Hochfeld-Magnetlabor Dresden, BEN YAGER, MATTHIAS ESCHRIG, JOHN SAUNDERS, Royal Holloway, University of London — Semimetals like graphite have recently received compelling interest as they not only are able to host topologically non-trivial phases but also can be driven into the ultra-quantum limit by magnetic fields now achievable in modern-day laboratories. Thus, they provide insight into quantum-Hall physics and the physics of massless Dirac fermions in three dimensions. They also represent ideal model systems for studying magnetic-field driven density wave instabilities, as the onset field for such collective excitations is suppressed in semimetals. Using pulsed high-magnetic fields up to 60 T applied to a single crystal of natural Tanzanian graphite, we find a series of field-induced phase transitions into collinear charge-density wave states resulting from enhanced interactions between the lowest four Landau levels. By analysing magneto-transport data and calculating the renormalized Landau level structure at high fields, we establish the phase diagram of graphite in its ultra-quantum limit. Our results imply the existence of a topologically-protected chiral edge state at high fields supporting both charge and spin currents.

¹We acknowledge the support of the HLD-HZDR, member of the European Magnetic Field Laboratory (EMFL), the Hubbard Theory Consortium, and the Engineering and Physical Science Research Council (EPSRC Grant Nos. EP/H048375/1 and EP/J010618/1)

Aldo Isidori
Royal Holloway, University of London

Date submitted: 05 Nov 2014

Electronic form version 1.4