

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Wire deconstructionism of two-dimensional topological phases¹

RONNY THOMALE, Wuerzburg University, TITUS NEUPERT, Princeton University, CLAUDIO CHAMON, Boston University, CHRISTOPHER MUDRY, PSI Zurich — A scheme is proposed to construct integer and fractional topological quantum states of fermions in two spatial dimensions. We devise models for such states by coupling wires of non-chiral Luttinger liquids of electrons, that are arranged in a periodic array. Which inter-wire couplings are allowed is dictated by symmetry and the compatibility criterion that they can simultaneously acquire a finite expectation value, opening a spectral gap between the ground state(s) and all excited states in the bulk. First, with these criteria at hand, we reproduce the tenfold classification table of integer topological insulators, where their stability against interactions becomes immediately transparent in the Luttinger liquid description. Second, we construct an example of a strongly interacting fermionic topological phase of matter with short-range entanglement that lies outside of the tenfold classification. Third, we expand the table to long-range entangled topological phases with intrinsic topological order and fractional excitations.

¹This work is supported by the European Research Council (ERC) through ERC-StG-Thomale-336012.

Ronny Thomale
Wuerzburg University

Date submitted: 12 Nov 2014

Electronic form version 1.4