

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Stabilization of CH₂ on Ru(0001) by hydrogen co-adsorption
SERGEY V. LEVCHENKO, XUNHUA ZHAO, MATTHIAS SCHEFFLER, Fritz-Haber-Institut der MPG, Berlin, FRITZ-HABER-INSTITUT DER MPG, BERLIN TEAM — Based on indirect experimental evidence, CH₂ was proposed as a building block for hydrocarbon chain growth on the Ru(0001) surface during the Fischer-Tropsch process. However, previous calculations agreed that CH₂ is not stable on Ru(0001) at the reaction conditions, and should quickly convert into CH. Employing density-functional theory, we show that this disagreement can be reconciled if co-adsorbed hydrogen is present on the surface. The atomic structure of various CH_x+H_y phases is obtained with genetic algorithm. CH_x dissociation barriers are calculated using the string method. We further demonstrate, by calculating the surface phase diagram for one-carbon species on Ru(0001) as a function of H₂ chemical potential, that the stabilization of CH₂ by co-adsorbed hydrogen requires non-equilibrium conditions. The calculated barrier for the CH₂ dissociation in the presence of hydrogen is significantly increased, and is close to the one recently measured by vibrational sum-frequency generation spectroscopy [1]. Our results also explain why CH₂ was not observed when C or CH are hydrogenated on Ru(0001), although it is observed after methane decomposition.

[1] H. Kirsch *et al.*, *J. Catal.* **320**, 89 (2014).

Sergey V. Levchenko
Fritz-Haber-Institut der MPG, Berlin

Date submitted: 14 Nov 2014

Electronic form version 1.4