

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Observation of Skyrmions in Mn-Pt-Sn based Heusler material

AJAYA KUMAR NAYAK, ROSHNEE SAHOO, Max Planck Institute for Chemical Physics of Solids, JOHANNES WILD, Faculty of the Institute of Experimental and Applied Physics, Regensburg, Germany, DANIEL EBKE, Max Planck Institute for Chemical Physics of Solids, JOSEF ZWECK, Faculty of the Institute of Experimental and Applied Physics, Regensburg, Germany, STUART PARKIN, Max Planck Institute of Microstructure Physics, CLAUDIA FELSER, Max Planck Institute for Chemical Physics of Solids — Mn₂YZ based Heusler materials often exhibit collinear spin alignment[1]. However, in a recent work we found Mn₂RhSn displays a non-collinear magnetic structure[2]. The presence of noncollinear spin alignment with lack of inversion symmetry sets up a perfect condition for skyrmions in Mn₂RhSn. However, experimentally it is still to be verified. To find skyrmions in Heusler system we search for materials with large spin-orbit coupling and lack of inversion symmetry. Here we present the experimental evidence of skyrmions in Mn-Pt-Sn Heusler compound. This sample shows an order-disorder transition around 400 K, followed by a spin-reorientation transition at low temperature. Our magnetization and ac-susceptibility measurements display a similar type of behavior that generally found in the skyrmion system. We have performed high resolution Lorentz transmission electron microscopy measurements to get a direct evidence of skyrmions in Mn-Pt-Sn sample. Our results show evolution of skyrmions in particular fields and temperature intervals.

[1] A. K. Nayak, et al., Phys. Rev. Lett. 110, 127204 (2013).

[2] O. Meshcheriakova, et al., Phys. Rev. Lett. 113, 08720(2014).

Claudia Felser
Max Planck Institute for Chemical Physics of Solids

Date submitted: 14 Nov 2014

Electronic form version 1.4