

Abstract Submitted  
for the MAR15 Meeting of  
The American Physical Society

**Pattern Recognition of Adsorbing HP Lattice Proteins<sup>1</sup>**

MATTHEW S. WILSON, GUANGJIE SHI, Center for Simulational Physics, University of Georgia, THOMAS WÜST, Scientific IT Services, ETH Zurich, DAVID P. LANDAU, Center for Simulational Physics, University of Georgia, FRIEDERIKE SCHMID, Institute of Physics, Johannes Gutenberg University — Protein adsorption is relevant in fields ranging from medicine to industry, and the qualitative behavior exhibited by course-grained models could shed insight for further research in such fields. Our study on the selective adsorption of lattice proteins utilizes the Wang-Landau algorithm to simulate the Hydrophobic-Polar (H-P) model<sup>2</sup> with an efficient set of Monte Carlo moves<sup>3</sup>. Each substrate is modeled as a square pattern of 9 lattice sites which attract either H or P monomers, and are located on an otherwise neutral surface. The fully enumerated set of 102 unique surfaces is simulated with each protein sequence. A collection of 27-monomer sequences<sup>4</sup> is used— each of which is non-degenerate and protein-like. Thermodynamic quantities such as the specific heat and free energy are calculated from the density of states, and are used to investigate the adsorption of lattice proteins on patterned substrates.

<sup>1</sup>Research supported by NSF.

<sup>2</sup>K. A. Dill, *Biochemistry* 24, 1501 (1985); K. F. Lau and K. A. Dill, *Macromolecules* 22, 3986 (1989).

<sup>3</sup>T. Wüst and D. P. Landau, *Phys. Rev. Lett.* 102, 178101 (2009); *J. Chem. Phys.* 137, 064903 (2012).

<sup>4</sup>M. Mann, D. Maticzka, R. Saunders, and R. Backofen. *HFSP Journal* 2. 396. Special issue on protein folding: experimental and theoretical approaches.(2008).

Matthew S. Wilson  
Center for Simulational Physics, University of Georgia

Date submitted: 14 Nov 2014

Electronic form version 1.4