

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Strong-coupling ansatz for the one-dimensional Fermi gas in a harmonic potential MEERA PARISH, London Centre for Nanotechnology, JESPER LEVINSEN, Aarhus University, PIETRO MASSIGNAN, ICFO, GEORG BRUUN, Aarhus University — The 1D Fermi gas with repulsive short-range interactions provides an important model of strong correlations and is often amenable to exact methods. However, in the presence of confinement, no exact solution is known for an arbitrary number of strongly interacting fermions. Here, we propose a novel ansatz for generating the lowest-energy wavefunctions of the repulsive 1D Fermi gas in a harmonic potential near the Tonks-Girardeau limit of infinite interactions. We specialize to the case of a single impurity interacting with N majority particles, where we may derive analytic forms of the approximate wavefunctions. Comparing with exact numerics, we show that the overlap between the wavefunctions from our ansatz and the exact ones in the ground-state manifold exceeds 0.9997 for $N \leq 8$. Moreover, the overlap for the ground-state wavefunction extrapolates to 0.9999 as $N \rightarrow \infty$. Thus our ansatz is essentially indistinguishable from numerically exact results in both the few- and many-body limits.

Meera Parish
London Centre for Nanotechnology

Date submitted: 14 Nov 2014

Electronic form version 1.4