

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Parallel Quantum Circuit in a Tunnel Junction¹ OMID FAIZY NAMARVAR, GHASSEN DRIDI, CHRISTIAN JOACHIM, PicoLab, CEMES-CNRS, GNS THEORY GROUP TEAM — In between 2 metallic nanopads, adding identical and independent electron transfer paths in parallel increases the electronic effective coupling between the 2 nanopads through the quantum circuit defined by those paths. Measuring this increase of effective coupling using the tunnelling current intensity can lead for example for 2 paths in parallel to the now standard $G = G_1 + G_2 + 2\sqrt{G_1 \cdot G_2}$ conductance superposition law (1). This is only valid for the tunnelling regime (2). For large electronic coupling to the nanopads (or at resonance), G can saturate and even decay as a function of the number of parallel paths added in the quantum circuit (3). We provide here the explanation of this phenomenon: the measurement of the effective Rabi oscillation frequency using the current intensity is constrained by the normalization principle of quantum mechanics. This limits the quantum conductance G for example to go when there is only one channel per metallic nanopads. This effect has important consequences for the design of Boolean logic gates at the atomic scale using atomic scale or intramolecular circuits.

References:

- (1) Joachim et al, Phys. Rev. B, 59, 16011 (1999).
- (2) C. Joachim, Nature Nano., 7, 620 (2012).
- (3) Sadeghi et al; PNAS 2015, 112, 9, 26

¹This has the financial support by European PAMS project

Omid Faizy Namarvar
CNRS/INSA

Date submitted: 23 Sep 2015

Electronic form version 1.4