

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Persistence **of**
**Dirac Node near Antiferromagnetic-to-Superconducting Phase Boundary
in $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$** HITOSHI TAKITA, NAOYA KISHIMOTO, YOUSUKE
NAKASHIMA, Hiroshima University, AKIHIRO INO, MASASHI ARITA, HIRO-
HUMI NAMATAME, MASAKI TANIGUCHI, Hiroshima Synchrotron Radiation
Center, YOSHIHIRO AIURA, IZUMI HASE, HIROSHI EISAKI, KUNIHIRO KI-
HOU, CHUL-HO LEE, AKIRA IYO, National Institute of Advanced Industrial
Science and Technology, MASAMICHI NAKAJIMA, Osaka University, SHIN-ICHI
UCHIDA, University of Tokyo, HIROSHIMA UNIVERSITY TEAM, HIROSHIMA
SYNCHROTRON RADIATION CENTER TEAM, NATIONAL INSTITUTE OF
INDUSTRIAL SCIENCE AND TECHNOLOGY TEAM, OSAKA UNIVERSITY
TEAM, UNIVERSITY OF TOKYO TEAM — Since the ground state of iron-
pnictides changes from an antiferromagnetic (AF) phase to a superconducting (SC)
phase, the evolution of electronic structure has attracted much attention. How-
ever, systematic investigation has been hindered by the intricate multiple bands
arising from the orbital degree of freedom of iron $3d$ states. Here we performed a
polarization-dependent ARPES study of $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ across the AF-SC phase
boundary. The doping-dependence of ARPES spectra has shown that the Dirac node
reported in the AF phase of BaFe_2As_2 persists in $x = 0.04$ near the AF-SC phase
boundary, and that it disappears in the SC phase of $x = 0.05$. We parametrized
the cone-like dispersion in $x = 0.04$. The polarization-dependence of our ARPES
spectra is consistent with the view that the Dirac node is protected by Berry phase
arising from orbital degree of freedom under the inversion symmetry.

Hitoshi Takita
Hiroshima University

Date submitted: 05 Nov 2015

Electronic form version 1.4