

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Nematic-columnar phase transition in oriented hard rectangles

TRISHA NATH, The Institute of Mathematical Sciences, C.I.T. Campus, Taramani, Chennai 600113, India, DEEPAK DHAR, Department of Theoretical Physics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai, 400005, India, R. RAJESH, The Institute of Mathematical Sciences, C.I.T. Campus, Taramani, Chennai 600113, India — We consider an assembly of monodispersed hard rectangles of size $2 \times d$ on a square lattice with only hard core interactions amongst them. The long axes of the rectangles can be oriented along the horizontal or vertical directions. For large enough aspect ratio, it is known that this system undergoes three phase transitions as the density (ρ) of rectangles is increased: first an isotropic-nematic transition (at ρ_1^*), second a nematic-columnar transition (at ρ_2^*), and third a columnar-sublattice transition (at ρ_3^*). In the nematic phase, only the orientational symmetry is broken. The columnar and sublattice phases correspond to additional broken translational symmetries along one (perpendicular to the nematic orientation) and both directions respectively. Interestingly, the critical value ρ_2^* remains finite, approximately 0.73, even as $d \rightarrow \infty$. We develop a systematic high density expansion for the surface tension between two differently-ordered columnar phases. Keeping only the first order perturbative correction term and setting this surface tension to zero, we get an estimate of ρ_2^* in excellent agreement with estimates from Monte Carlo simulations, for all $d \geq 2$.

Trisha Nath
The Institute of Mathematical Sciences, C.I.T. Campus, Taramani, Chennai 600113, India

Date submitted: 05 Nov 2015

Electronic form version 1.4