

Abstract Submitted  
for the MAR16 Meeting of  
The American Physical Society

**Statistical Transmutation in Periodically Driven Optical Lattices<sup>1</sup>**

TIGRAN SEDRAKYAN, W. I. Fine Theoretical Physics Institute, University of Minnesota, Joint Quantum Institute, University of Maryland - College Park, VICTOR GALITSKI, Joint Quantum Institute, University of Maryland - College Park, ALEX KAMENEV, W. I. Fine Theoretical Physics Institute, University of Minnesota — We show that interacting bosons in a periodically driven two dimensional (2D) optical lattice may effectively exhibit fermionic statistics. The phenomenon is similar to the celebrated Tonks-Girardeau regime in 1D. The Floquet band of a driven lattice develops the moat shape, i.e., a minimum along a closed contour in the Brillouin zone. Such degeneracy of the kinetic energy favors fermionic quasiparticles. The statistical transmutation is achieved by the Chern-Simons flux attachment similar to the fractional quantum Hall case. We show that the velocity distribution of the released bosons is a sensitive probe of the fermionic nature of their stationary Floquet state.

<sup>1</sup>This work was supported by the PFC-JQI (T.S.), USARO and Simons Foundation (V.G.), and DOE contract DE-FG02-08ER46482 (A.K.)

Tigran Sedrakyan  
Univ of Minn - Minneapolis

Date submitted: 06 Nov 2015

Electronic form version 1.4