

Abstract Submitted  
for the MAR16 Meeting of  
The American Physical Society

**Unified boundary conditions and Casimir forces for fields with arbitrary spin**<sup>1</sup> ROBERT BENNETT, University of Freiburg, ADAM STOKES, University of Leeds — The electromagnetic Casimir effect is well-known and has been extensively studied for the last half-century. This attractive force between parallel plates arises from the imposition of boundary conditions upon the fluctuating spin-1 photon field, so a natural further question is whether fields of different spin can cause similar forces when confined in the same way. However, so far it has not been clear what the appropriate boundary conditions for physically-confined spinor fields may be. Here we present work that generalises the physically well-motivated electromagnetic boundary conditions to fields of arbitrary spin, thus arriving at physically reasonable boundary conditions<sup>2</sup> and Casimir forces<sup>3</sup> for a selection of interesting fields. For example, the so-called ‘bag model’ boundary conditions from nuclear physics emerge from our generalised boundary condition as a special case, as do the linearised gravity boundary conditions suggested in a remarkable recent proposal<sup>4</sup> concerning possible measurement of gravitonic Casimir forces.

<sup>1</sup>Supported by the UK Engineering and Physical Sciences Research Council (EPSRC)

<sup>2</sup>A. Stokes and R. Bennett, *New J. Phys.* 17 073012 (2015)

<sup>3</sup>A. Stokes and R. Bennett, *Ann. Phys NY* 360 246 (2015)

<sup>4</sup>James Q. Quach, *Phys. Rev. Lett.* 114 081104 (2015)

Robert Bennett  
University of Freiburg

Date submitted: 06 Nov 2015

Electronic form version 1.4