

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Effect of longer-range lattice anisotropy on the electronic structure and magnetism of spin-orbit-coupled 5d transition-metal oxides
NIKOLAY BOGDANOV¹, VAMSHI KATUKURI², JUDIT ROMHÁNYI³, Institute for Theoretical Solid State Physics, IFW Dresden, Germany, VIKTOR YUSHANKHAI, Joint Institute for Nuclear Research, Dubna, Russia, VLADISLAV KATAEV, BERNDT BÜCHNER, Institute for Solid State Research, IFW Dresden, Germany, JEROEN VAN DEN BRINK, LIVIU HOZOI, Institute for Theoretical Solid State Physics, IFW Dresden, Germany — Our detailed quantum chemistry calculations provide magnetic g factors and exchange interactions for the quasi two-dimensional iridates Sr_2IrO_4 and Ba_2IrO_4 . While canonical ligand-field considerations predict g_{\parallel} -factors < 2 for the positive tetragonal distortions present in Sr_2IrO_4 , we find $g_{\parallel} > 2$. This implies that the d levels in Sr_2IrO_4 are inverted with respect to the ordering deduced from the local ligand distortions, whereas we find them in Ba_2IrO_4 to be instead normally ordered. Electron spin resonance measurements confirm the level inversion in Sr_2IrO_4 . This d -level switching is driven by the specific ionic charge distribution within adjacent IrO_2 and SrO layers. Since polar discontinuities and the associated complications do not arise for such layers, our results highlight the tetravalent d -metal 214 oxides as ideal platforms to explore d -level reconstruction and engineering in the context of oxide heterostructures.

¹Present address: Electronic structure theory, MPI-FKF, Stuttgart

²Present address: Ecole polytechnique fédérale de Lausanne

³Present address: Quantum materials, MPI-FKF, Stuttgart

Nikolay Bogdanov
Institute for Theoretical Solid State Physics, IFW Dresden

Date submitted: 06 Nov 2015

Electronic form version 1.4