

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Spatially Mapping Dirac Fermions in a Graphene Quantum Dot

JUWON LEE, DILLON WONG, JAIRO VELASCO JR, Univ of California - Berkeley, JOAQUIN RODRIGUEZ-NIEVA, MIT, SALMAN KAHN, Univ of California - Berkeley, PHONG VO, MIT, HSIN-ZON TSAI, Univ of California - Berkeley, TAKASHI TANIGUCHI, KENJI WATANABE, National Institute for Materials Science, ALEX ZETTL, FENG WANG, Univ of California - Berkeley, LEONID LEVITOV, MIT, MICHAEL CROMMIE, Univ of California - Berkeley — Quantum confinement in graphene is important for tuning and exploiting graphene's electronic, spin, and optical properties. Here we present a novel technique for creating gate-tunable graphene quantum dots that are fully exposed and compatible with surface characterization tools. Using scanning tunneling microscopy (STM), we are able to spatially visualize and characterize the electronic structure of these gate-tunable graphene quantum dots. The quantum interference patterns observed in this way can be compared to the predictions of the Dirac equation, thus providing new insight into the behavior of confined ultra-relativistic Dirac fermions.

Juwon Lee
Univ of California - Berkeley

Date submitted: 06 Nov 2015

Electronic form version 1.4