

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

A Tight Binding Approach to Understanding the Orbital Character of Pb-BSSCO and its Effects on Photoemission Matrix Elements¹

THOMAS NUMMY, University of Colorado Boulder, YUE CAO, Brookhaven National Lab, JUSTIN WAUGH, STEPHEN PARHAM, HAOXIANG LI, XIAOQUING ZHOU, University of Colorado Boulder, SUNG-KWAN MO, Lawrence Berkeley National Lab, A. MAGREZ, H. BERGER, EPFL Lausanne, DANIEL DESSAU, University of Colorado Boulder — We measure the near Fermi energy electronic structure of Pb-BSSCO using Angle Resolved Photoemission Spectroscopy (ARPES) for various light polarization, incident at a glancing angle. A strong dependence of the photocurrent for a given crystal momentum on the polarization of light is observed. A minimal model tight binding calculation in a basis of local atomic orbitals is then utilized to determine the orbital composition of the single particle wavefunction as a function of crystal momentum in a single CuO₂ plane. Building off of these results, we simulate the relative ARPES photocurrent throughout crystal momentum space in the dipole approximation and compare to our experimental data. These comparisons confirm the presence of changing wavefunction orbital composition throughout the Brillouin zone in Pb-BSSCO.

¹Acknowledgement: NSF GRFP

Tom Nummy
University of Colorado Boulder

Date submitted: 06 Nov 2015

Electronic form version 1.4