

Abstract Submitted
for the MAR17 Meeting of
The American Physical Society

Proton radiography measurements of ejecta structure in shocked Sn¹ J.E. HAMMERBERG, W.T. BUTTLER, A. LLOBET, C. MORRIS, Los Alamos National Laboratory — We have performed ejecta measurements at the Los Alamos proton radiography facility on 7 mm thick 50 mm diameter Sn samples driven with a PBX9501 high explosive. The surface of the Sn, in contact with He gas at an initial pressure of 7 atmospheres, was machined to have 3 concentric sinusoidal features with a wavelength of $\lambda = 2\text{mm}$ in the radial direction and amplitude $h_0 = 0.159\text{mm}$ ($kh_0 = 2\pi h_0/\lambda = 0.5$). The shock pressure was 27 GPa. 28 images were obtained between 0 and 14 μs from the time of shock breakout at 500 ns intervals. The Abel inverted density profiles evolve to a self-similar density distribution that depends on a scaling variable $z/v_s t$ where v_s is the spike tip velocity, z is the distance from the free surface and t is the time after shock breakout. Both the density profiles and the time dependence of the mass per unit area in the evolving spikes are in good agreement with a Richtmyer-Meshkov instability based model for ejecta production and evolution.

¹This work was performed under the auspices of the U.S. Dept. of Energy under contract DE-AC52-06NA25396. The support of the LANL ASC- PEM and Science Campaign 2 programs is gratefully acknowledged.

J.E. Hammerberg
Los Alamos National Laboratory

Date submitted: 20 Nov 2016

Electronic form version 1.4