

Abstract Submitted
for the MAR17 Meeting of
The American Physical Society

Tuning spin-polarized transport in organic semiconductors

RICHARD MATTANA, MARTA GALBIATI, SOPHIE DELPRAT, SERGIO TATAY, CYRILE DERANLOT, PIERRE SENEOR, FREDERIC PETROFF, Unit Mixte de Physique CNRS/Thales — Molecular spintronics is an emerging research field at the frontier between organic chemistry and the spintronics. Compared to traditional inorganic materials molecules are flexible and can be easily tailored by chemical synthesis. Due to their theoretically expected very long spin lifetime, they were first only seen as the ultimate media for spintronics devices. It was recently that new spintronics tailoring could arise from the chemical versatility brought by molecules. The hybridization between a ferromagnet and molecules induces a spin dependent broadening and energy shifting of the molecular orbitals leading to an induced spin polarization on the first molecular layer. This spin dependent hybridization can be used to tailor the spin dependent transport in organic spintronics devices. We have studied vertical Co/Alq3/Co organic spin valves. The negative magnetoresistance observed is the signature of different coupling strengths at the top and bottom interfaces. We have then inserted an inorganic tunnel barrier at the bottom interface in order to suppress the spin-dependent hybridization. In this case we restore a positive magnetoresistance. This demonstrates that at the bottom Co/Alq3 interface a stronger coupling occurs which induces an inversion of the spin polarization.

Richard Mattana
Unit Mixte de Physique CNRS/Thales

Date submitted: 10 Nov 2016

Electronic form version 1.4