

Abstract Submitted  
for the MAR17 Meeting of  
The American Physical Society

**Conditions for  $T^2$  resistivity from electron-electron scattering<sup>1</sup>**

MICHAEL SWIFT, CHRIS G. VAN DE WALLE, Univ of California - Santa Barbara — Many complex oxides (including titanates, nickelates and cuprates) exhibit a carrier scattering mechanism with a power law dependence on temperature (resistivity  $\rho \propto T^2$ ). By analogy to a similar phenomenon observed in metals at low temperature, this mechanism has often been identified as Fermi-liquid-like electron-electron scattering (Baber scattering). However, other transport signatures in these materials have shown behavior that casts doubt on this simple picture. Careful investigation of electron-electron scattering (both direct and phonon-mediated) reveals that the Baber  $T^2$  power law rests on several crucial assumptions. In some cases, these assumptions are not satisfied and removing them destroys the power law. We illustrate these issues with two case studies: sodium metal (in which electron-electron scattering gives  $T^2$  resistivity) and strontium titanate (in which it does not). Our results suggest that an observation of  $\rho \propto T^2$  is not sufficient evidence for electron-electron scattering. The power law observed in the complex oxides may instead be due to another, as yet undiscovered, mechanism.

<sup>1</sup>This work was supported by ONR.

Michael Swift  
Univ of California - Santa Barbara

Date submitted: 10 Nov 2016

Electronic form version 1.4