

Abstract Submitted
for the MAR17 Meeting of
The American Physical Society

Monte Carlo Simulation of Amyloid Protofibril Formation

MATTHEW S. WILSON, GUANGJIE SHI, Center for Simulational Physics, University of Georgia, THOMAS WÜST, Scientific IT Services, ETH Zurich, DAVID P. LANDAU, Center for Simulational Physics, University of Georgia, FRIEDERIKE SCHMID, Institute of Physics, Johannes Gutenberg University — Aggregation processes of amyloid protofibrils are studied using the replica-exchange Wang-Landau (REWL)¹ algorithm to simulate multiple interacting model peptides. The HOP model², which adds an additional neutral polarity group to the classic hydrophobic-polar (HP) model³, is used for simplicity and efficiency. Constituent peptides are modeled as short, intrinsically disordered HOP sequences, which do not form globular structures individually but self-assemble to form various aggregated structures. Using the parallelized sampling framework, the density of states is determined and the minimal energy state is identified. From thermodynamic quantities, the effects of peptide concentration are studied for the formation of protofibrillar structures.

¹T. Vogel, Y. W. Li, T. Wüst, and D. P. Landau, Phys. Rev. E **90**, 023302 (2014).

²G. Shi, T. Wüst, Y. W. Li, and D. P. Landau, J. Phys.: Conf. Ser. **640**, 012017 (2015).

³K. A. Dill, Biochemistry **24**, 1501 (1985); K. F. Lau and K. A. Dill, Macromolecules **22**, 3986 (1989).

Matthew S. Wilson
Center for Simulational Physics, University of Georgia

Date submitted: 10 Nov 2016

Electronic form version 1.4