

Abstract Submitted
for the APR05 Meeting of
The American Physical Society

Sorting Category: F8. (E)

Measuring Active-Sterile Neutrino Oscillations with a Stopped Pion Neutrino Source RICHARD VAN DE WATER, GORDON MCGREGOR, Los Alamos National Laboratory — The question of the existence of light sterile neutrinos is of great interest in many areas of particle physics, astrophysics, and cosmology. Furthermore, should the MiniBooNE experiment at Fermilab confirm the LSND oscillation signal, then new measurements are required to identify the mechanism responsible for these oscillations. Possibilities include sterile neutrinos, CP or CPT violation, variable mass neutrinos, and Lorentz violation. In this paper, we consider an experiment at a stopped pion neutrino source to determine if active-sterile neutrino oscillations with Δm^2 greater than 0.1 eV^2 can account for the signal. By exploiting stopped π^+ decay to produce a monoenergetic ν_μ source, and measuring the rate of the neutral current reaction $\nu_x \text{}^{12}\text{C} \rightarrow \nu_x \text{}^{12}\text{C}^*$ (15.11) as a function of distance from the source, we show that a convincing test for active-sterile neutrino oscillations can be performed.

Prefer Oral Session
 Prefer Poster Session

Richard Van de Water
vdwater@lanl.gov
Los Alamos National Laboratory

Date submitted: 13 Jan 2005

Electronic form version 1.4