

Abstract Submitted
for the DAMOP05 Meeting of
The American Physical Society

Sorting Category: 3. (T)

Semiclassical treatment of $\bar{p}p$ formation in \bar{p} -H collisions R. CABRERA-TRUJILLO, B.D. ESRY, J. R. Macdonald Laboratory, Department of Physics, Kansas State University, Manhattan, KS 66506 — As the energy of an antiproton colliding with a hydrogen atom decreases, the probability for formation of protonium ($\bar{p}p$) increases. In this work, we present a calculation of protonium formation using the Electron-Nuclear Dynamics (END)¹ theory for projectile energies from 1 eV to 10 eV. We present preliminary results for the protonium formation cross section, the stopping cross section (nuclear and electronic). In particular, we explore the role of non-adiabatic effects and the ionization channel within the END formalism.

¹E. Deumens, A. Diz, R. Longo, and Y. Öhrn, Rev. Mod. Phys. **66**, 917 (1994).

- Prefer Oral Session
 Prefer Poster Session

Remigio Cabrera-Trujillo
trujillo@phys.ksu.edu
J. R. Macdonald Laboratory, Department of Physics
Kansas State University, Manhattan, KS 66506

Date submitted: 01 Feb 2005

Electronic form version 1.4