

Abstract Submitted
for the DAMOP05 Meeting of
The American Physical Society

Sorting Category: 2.4 (E)

Probing Vibrational and Dissociative Wavepacket Dynamics in H₂ and D₂ Molecules with Ultra Short Laser Pulses¹

ALI ALNASER, B. ULRICH, X.-M. TONG, C.M. MAHARJAN, P. RANITOVIC, T. OSIPOV, R. ALI, B. SHAN, Z. CHANG, C.D. LIN, I.V. LITVINYUK, C.L. COCKE, J. R. MACDONALD LABORATORY, PHYSICS DEPARTMENT, KANSAS STATE UNIV., MANHATTAN, KS, 66506 TEAM — We have used ultra short laser pulses of 8fs duration in pump- probe arrangement to probe fast wavepacket dynamics in H₂ (D₂) molecules. A vibrational wave packet is launched onto the $1s\sigma_g$ potential curve of the singly charged molecule by the (tunneling) removal of one electron from the neutral molecule. By choosing the proper laser intensities for the pump and the probe pulses, we were able to monitor the simultaneous propagation of this wave packet on the (coupled) $1s\sigma_g$ (bound) and $2p\sigma_u$ (dissociative) potential curves of H₂⁺ and D₂⁺ over a time delay of 0-100 fs between the two pulses. We used the COLTRIMS technique to measure the kinetic energy release (KER) of the proton (deuteron) pairs produced in the Coulomb explosion of these molecules as function of time. A quantitative model has been constructed which accounts for the major features observed in the experiment.

¹This work was supported by Chemical Sciences, Geosciences and Biosciences Division, Office of Basic Energy Sciences, Office of Science, U. S. Department of Energy

Prefer Oral Session
 Prefer Poster Session

Ali Alnaser
alnaser@phys.ksu.edu

Date submitted: 28 Jan 2005

Electronic form version 1.4