

Abstract Submitted
for the DAMOP05 Meeting of
The American Physical Society

Sorting Category: 2.3 (E)

Parallel and perpendicular transitions in the dissociation of O_2^+ caused by an ultrashort intense laser pulse¹ A.M. SAYLER, P.Q. WANG, K.D. CARNES, B.D. ESRY, I. BEN-ITZHAK, J.R. Macdonald Laboratory, Department of Physics, Kansas State University — Laser-induced dissociation of O_2^+ has been experimentally studied using ultrashort (~ 50 fs) intense ($\sim 10^{14}$ W/cm²) laser pulses at 790 nm. The O^+ and O fragments are measured in coincidence with a 3-dimensional momentum imaging system, which provides both angular and kinetic energy release (KER) distributions for dissociation. By analyzing the angular distribution of a specific range in the KER spectrum, the dissociation pathway may be deduced. In particular, a single photon parallel transition ($\Delta\Lambda=0$) is expected to have a $\cos^2\theta$ distribution, while a perpendicular one ($\Delta\Lambda=1$) leads to a $\sin^2\theta$ distribution. Therefore, a dissociation pathway requiring the exchange of n $\Delta\Lambda=0$ photons and m $\Delta\Lambda=1$ photons is expected to have a $\cos^{2n}\theta\sin^{2m}\theta$ distribution. Thus, knowing n and m along with the KER range allows a plausible identification of the dissociation pathway.

¹Supported by the Chemical Sciences, Geosciences and Biosciences Division, Office of Basic Energy Sciences, Office of Science, U.S. Department of Energy.

- Prefer Oral Session
 Prefer Poster Session

J.R. Macdonald Laboratory, Department of Physics, Kansas State University

A.M. Sayler
sayler@phys.ksu.edu

Date submitted: 01 Feb 2005

Electronic form version 1.4