

Abstract Submitted  
for the DAMOP06 Meeting of  
The American Physical Society

Sorting Category: 2.1 (T)

**Term dependence of nondipole valence s-photoelectron angular distributions from half-filled shell atoms<sup>1</sup>**

VALERIY DOLMATOV, EMRE GULER<sup>2</sup>, University of North Alabama, Florence, AL, STEVEN MANSON, Georgia State University, Atlanta, GA — To date, nondipole photoelectron angular distributions from open-shell atoms have remained poorly studied because of their complexity in both experimental and theoretical studies. To remedy this situation, we have initiated a theoretical research program for studying nondipole effects in these atoms. The aim of this presentation is to demonstrate a dramatic dependence of nondipole valence s-photoelectron angular distributions on final state terms of the open-shell atom-remainder. It turns out that electron correlation affects these nondipole angular distributions crucially but much differently depending on a final state term of the residual ion. As a result, there are marked differences between nondipole contributions to the valence s-photoelectron angular distributions “belonging” to different final state terms of the ion-remainder. This is demonstrated by our calculations for the 4s photoelectrons from the Mn( $3d^5 4s^2, ^6S$ ) atom for the  $4s, ^5S$  and  $4s, ^7S$  final state terms of the ion-remainder. Calculations were performed in a “spin-polarized” random-phase-approximation with exchange (SP RPAE) framework.

<sup>1</sup>The NSF and NASA financial support is acknowledged.

<sup>2</sup>Undergraduate student

Prefer Oral Session  
 Prefer Poster Session

Valeriy Dolmatov  
vkdolmatov@una.edu  
University of North Alabama, Florence, AL

Date submitted: 25 Jan 2006

Electronic form version 1.4