

Abstract Submitted
for the DFD07 Meeting of
The American Physical Society

Sorting Category: 22. (E)

The 3D flow structure of puffs in transitional pipe flow

JERRY WESTERWEEL, Delft University of Technology, CASIMIR VAN DOORNE, DIRK-JAN KUIK — Time-resolved stereo-PIV measurements were used to study the 3D flow structure of a puff in a pipe. At the trailing edge of the puff, where the laminar flow undergoes a transition to turbulence, pairs of counter rotating streamwise vortices result in large mushroom-like structures as seen in the LIF flow visualizations. The stereo-PIV system is used to take time-resolved measurements over the entire cross-section of the pipe. When time is converted to a spatial coordinate (assuming Taylor's hypothesis of 'frozen turbulence') we obtain the quasi-instantaneous flow structure of a turbulent puff. At the upstream end of the puff a quasi-periodic regeneration of streamwise vortices takes place. Initially the vortex structure resembles a travelling wave solution, but as the vortices propagate further into the turbulent region of the puff they continue to develop into strong hairpin vortices. The structure suggests a mechanism for the long persistence of these puffs.

- Prefer Oral Session
 Prefer Poster Session

Jerry Westerweel
j.westerweel@tudelft.nl
Delft University of Technology

Date submitted: 03 Aug 2007

Electronic form version 1.4