

Abstract Submitted
for the DFD07 Meeting of
The American Physical Society

Sorting Category: 11. (T)

Memory and inertial oscillations in bubble break-up

LAURA E. SCHMIDT, WENDY W. ZHANG, University of Chicago

— The break-up of a submerged air bubble has been observed to exhibit an exceptional singularity dynamics, one that retains a memory of small asymmetries in the initial shape [Keim et al. PRL **97**, 144503 (2006)]. To understand the origin of this memory, we observe that the slender-body model of cylindrically symmetric break-up dynamics has a Hamiltonian structure. It therefore retains a precise memory of the energy distribution at the onset of break-up. Moreover, our analysis shows this memory is preserved when the dynamics is perturbed from cylindrical symmetry. The initial asymmetry simply excites inertial oscillations of approximately constant amplitude about the $O(1)$ radial collapse. Finally, to connect our results with shape oscillations that have been recently observed in experiments [Keim & Nagel, DFD07], we include effects of surface tension. Surface tension effects are significant just after onset and act to speed up the oscillation and increase its amplitude. However, as the break-up singularity approaches, the growth of the inertial oscillation amplitude due to surface tension becomes negligible. Surface tension merely distorts the initial energy distribution, which is then remembered by the singularity.

Prefer Oral Session
 Prefer Poster Session

Laura E. Schmidt
schmidt@uchicago.edu
University of Chicago

Date submitted: 03 Aug 2007

Electronic form version 1.4