

Abstract Submitted
for the DNP07 Meeting of
The American Physical Society

Sorting Category: 1. (E)

Determination of the astrophysical S-factor for the $^{12}\text{N}(\text{p},\gamma)^{13}\text{O}$ reaction from ($^{12}\text{N},^{13}\text{O}$) proton transfer reaction
BANU, T. AL-ABDULLAH, C. FU, C.A. GAGLIARDI, Y. LI, M. MCCLESKEY, G. TABACARU, L. TRACHE, R.E. TRIBBLE, Y. ZHAI, Cyclotron Institute, Texas A&M University, College Station, TX 77843, V. BURJAN, Institute of Nuclear Physics, Czech Academy of Sciences, Prague, Czech Republic, F. CARSTOIU, IFIN-HH, Bucharest, Romania — The reaction rate for the radiative proton capture on the drip line nucleus ^{12}N was determined using the indirect Asymptotic Normalization Coefficient (ANC) method. This reaction is important for studying the nucleosynthesis in Population III stars with low-metallicity. A 23 MeV/nucleon ^{12}C primary beam from the K500 cyclotron at Texas A&M University was employed. Secondary ^{12}N beam of 2×10^5 pps was separated using the recoil spectrometer MARS. The $^{14}\text{N}({}^{12}\text{N},{}^{13}\text{O})^{13}\text{C}$ proton transfer reaction at 12 MeV/nucleon was measured to extract the ANC for the virtual decay $^{13}\text{O} \rightarrow ^{12}\text{N} + \text{p}$. The ANC was then used to determine the direct component of the astrophysical S-factor. The results of this measurement will be discussed.

- Prefer Oral Session
 Prefer Poster Session

Adriana Banu
banu@comp.tamu.edu
Cyclotron Institute, Texas A&M University, College Station, TX 77843

Date submitted: 09 Aug 2007

Electronic form version 1.4