

Abstract Submitted
for the DNP07 Meeting of
The American Physical Society

Sorting Category: 1. (T)

GFMC Calculations of Isospin-Mixing in ^8Be ¹ ROBERT B. WIRINGA, STEVEN C. PIEPER, MUSLEMA PERVIN, Argonne National Laboratory — ^8Be has two 2^+ states at 16.6 and 16.9 MeV excitation that are strongly isospin-mixed. We have performed microscopic Green's function Monte Carlo calculations of the isospin-mixing matrix elements between the T=0 and 1 states using the realistic Argonne v_{18} + Illinois-2 Hamiltonian, which includes strong CSB components and a full electromagnetic interaction. We obtain 80% of the empirically-determined matrix element, with 2/5 of our result coming from the terms beyond Coulomb, confirming an earlier variational study. We have also calculated the mixing between the nearby T=0,1 pairs of 1^+ and 3^+ states. Finally, we have examined the mixing of the T=1 2^+ states with the first T=0 excited 2^+ state at 3.0 MeV, which is the final state for weak decays from either ^8Li or ^8B . We find this state, which is an important laboratory for testing various aspects of weak interactions, to have extremely small T=1 contamination.

¹Work supported by U.S. Department of Energy, Office of Nuclear Physics, under Contract No. DE-AC02CH11357; computations performed at the LCRC facility of Argonne National Laboratory

- Prefer Oral Session
 Prefer Poster Session

Robert Wiringa
wiringa@anl.gov
Argonne National Laboratory

Date submitted: 02 Jul 2007

Electronic form version 1.4