

Abstract Submitted  
for the DNP07 Meeting of  
The American Physical Society

Sorting Category: 1. (E)

**Photodisintegration Cross Section Measurements for  
 $^{142}\text{Nd}$  and  $^{150}\text{Nd}$  and Low-energy E1  $\gamma$ -ray Strength Functions**

C.T. ANGELL, UNC/TUNL, H. UTSUNOMIYA, S. GOKO, A. MAKINAGA, T. KAIHORI, Konan U., H. TOYOKAWA, AIST, Y.W. LUI, Texas A&M — The photo-neutron disintegration cross sections were measured near threshold for  $^{142}\text{Nd}$ , and, for the first time,  $^{150}\text{Nd}$ . The measurements were made using the monoenergetic  $\gamma$ -ray beam at the AIST TERAS facility in Tsukuba, Japan. The  $\gamma$ -ray beam was produced via laser inverse-Compton scattering. The neutrons were detected using a composite detector consisting of  $^3\text{He}$  proportional counters embedded in a polyethylene block. The technique and facility will be overviewed, highlighting the advantages of a monoenergetic  $\gamma$ -ray beam to absolute cross section measurements. The results will be compared to theoretical calculations using the Quasi-particle Random Phase Approximation (QRPA). Finally, the neutron number dependence of the low-energy E1  $\gamma$ -ray strength function will be discussed.

- Prefer Oral Session  
 Prefer Poster Session

Christopher Angell  
cangell@physics.unc.edu  
UNC/TUNL

Date submitted: 03 Jul 2007

Electronic form version 1.4