

Abstract Submitted
for the DNP07 Meeting of
The American Physical Society

Sorting Category: 11. (T)

Effective Interactions for pf Shell from the No Core Shell Model¹

ERDAL DIKMEN, Suleyman Demirel University, The University of Arizona, ALEXANDER LISETSKY, BRUCE BARRETT, The University of Arizona, PETR NAVÁTIL, Lawrence Livermore National Laboratory, JAMES VARY, Iowa State University — Following the ideas of the Tucson nuclear theory group for the extension of the No-Core Shell Model (NCSM) approach to sd-shell nuclei [1], we show how a NCSM calculation for ^{42}Ca in a $2\hbar\Omega$ model space can yield two-body effective interactions for the pf-shell. We demonstrate how the effective Hamiltonian derived in the $2\hbar\Omega$ NCSM at the 2-body cluster level should be modified to properly account for the many-body correlations produced by truncating to a single major shell. The pf-shell two-body effective interactions for ^{42}Ca , obtained by direct projection, are used to reproduce the results of large scale NCSM for other Ca isotopes.

1. B.R. Barrett et al, DNP 2007 abstract.

¹E.D. supported partly by TUBITAK-BIDEB and TUBITAK 105T092. B.R.B. and A.F.L. supported partly by NSF grant PHY-0555396. Work by P.N. partly performed under DOE contract No. W-7405-Eng-48. J.P.V. supported partly by USDOE grant DE-FG-02-87ER40371.

Prefer Oral Session
 Prefer Poster Session

Erdal Dikmen
edikmen@fef.sdu.edu.tr
Suleyman Demirel University, University of Arizona

Date submitted: 02 Jul 2007

Electronic form version 1.4