

Abstract Submitted  
for the DNP08 Meeting of  
The American Physical Society

Sorting Category: 6. (E)

**Isotopic width distributions and symmetry energy**

SARA WUENSCHEL, STRATOS GALANOPOULOS, KRIS HAGEL, ZACH KOHLEY, DINESH SHETTY, SARAH SOISSON, GEORGE SOULIOTIS, BRIAN STEIN, SHERRY YENNELLO, Texas A&M University Cyclotron Institute — Within the Microcanonical Multifragmentation Model (MMM), the bulk versus surface contribution to the symmetry energy is predicted to be distinguishable. If the symmetry energy is dominated by the surface, the symmetry energy coefficient should evolve with the size of the fragment studied. However, if the bulk term dominates, the symmetry energy coefficient should be constant across all fragment sizes. Symmetry energy can be accessed through knowledge of the parameter  $\alpha$ . This parameter may be obtained by isoscaling [1] or isotopic widths [2]. Differentiation between bulk and surface requires isotopic widths across a wide array of elements. Projectile fragmentation of  $^{86}\text{Kr}+^{64}\text{Ni}$  at 35MeV/u was taken on the NIMROD-ISiS detector. The wide range of isotopic resolution seen in NIMROD-ISiS data allows the isotopic widths for  $Z=3-15$  to be extracted for this study. Events are characterized through reconstruction of the quasi-projectile. Isotopic width data will be presented.

- Prefer Oral Session  
 Prefer Poster Session

Sara Wuenschel  
wuenschel@comp.tamu.edu  
Texas A&M University Cyclotron Institute

Date submitted: 02 Jul 2008

Electronic form version 1.4