

Abstract Submitted
for the DPP06 Meeting of
The American Physical Society

Sorting Category: 4.1.2 (C)

**Collisional Relaxation of Super Thermal Electrons
Generated by Relativistic Laser Pulses in Thin Solid Targets¹**

ANDREAS KEMP, EMMANUEL D'HUMIERES, YASUHIKO SENTOKU, University of Nevada, Reno, HUI CHEN, HYUN CHUNG, STEPHANIE HANSEN, RONNIE SHEPHERD, SCOTT WILKS, LLNL — The interaction of intense fs laser pulses with matter generates energetic electrons that penetrate deeply into solid targets and deposit energy on sub-ps time scales. Experiments at LLNL's Comet laser facility have used a ps time-resolved K-alpha diagnostic for hot electrons, allowing us to study the highly transient energy transfer between hot and thermal electrons. For hot populations with temperatures of ~ 500 keV generated by 0.5 ps laser pulses interacting with $12.5 \mu\text{m}$ thick Titanium slabs, relaxation times were found to be of the order of 20 ps [1]. Motivated by the discrepancy between the laser- and the K-alpha time scales, we study the roles of various effects that determine the generation of K-alpha radiation: collisional coupling between hot and thermal electrons, plasma expansion, and ionization. [1] H.Chen et al, this conference

¹Work supported by DOE/NNSA-UNR grant DE-FC52-01NV14050.

Prefer Oral Session
 Prefer Poster Session

Andreas Kemp
kemp@physics.unr.edu
University of Nevada, Reno

Date submitted: 31 Jul 2006

Electronic form version 1.4