

Abstract Submitted  
for the DPP07 Meeting of  
The American Physical Society

Sorting Category: 1.0.0 (T)

**Breathing Mode in Complex Plasmas**<sup>1</sup> K. FUJIOKA, City College of New York, C. HENNING, P. LUDWIG, M. BONITZ, ITAP, Christian-Albrechts-Universitaet zu Kiel, A. MELZER, Institute of Physics, Greifswald University, S. VITKALOV, City College of New York — The breathing mode is a fundamental normal mode present in Coulomb systems, and may have utility in identifying particle charge and the Debye length of certain systems. The question remains whether this mode can be extended to strongly coupled Yukawa balls [1]. These systems are characterized by particles confined within a parabolic potential well and interacting through a shielded Coulomb potential [2,3]. The breathing modes for a variety of systems in 1, 2, and 3 dimensions are computed by solving the eigenvalue problem given by the dynamical (Hesse) matrix. These results are compared to theoretical investigations that assume a strict definition for a breathing mode within the system, and an analysis is made of the most fitting model to utilize in the study of particular systems of complex plasmas [1,4].

References

- [1] T.E. Sheridan, Phys. of Plasmas. 13, 022106 (2006)
- [2] C. Henning et al., Phys. Rev. E 74, 056403 (2006)
- [3] M. Bonitz et al., Phys. Rev. Lett. 96, 075001 (2006)
- [4] C. Henning et al., submitted for publication

<sup>1</sup>Supported by the DFG (via SFB-TR24) and the DAAD RISE Program

- Prefer Oral Session
- Prefer Poster Session

Kenji Fujioka  
kenji.fujioka@gmail.com  
City College of the City University of New York

Date submitted: 20 Jul 2007

Electronic form version 1.4