

Abstract Submitted
for the MAR05 Meeting of
The American Physical Society

Sorting Category: 08.4 (C)

Oriental Anisotropy and Effective Electrostatic Interactions in Polyelectrolyte Star Solutions¹ HAO WANG, ALAN R. DENTON, Department of Physics, North Dakota State University — In solutions of polyelectrolyte (PE) star macroions, electrostatic interactions between charged arms of approaching macroions can induce anisotropy in the arm orientational distributions. Within a rigid arm model, we explore the influence of arm orientational anisotropy on the effective pair interactions between stars. For strongly charged arms, where electrostatic energies much exceed thermal energies, a torque balance calculation predicts significant arm anisotropy and a corresponding change in the effective pair interaction between stars. For weakly charged arms where electrostatic and thermal energies are comparable, two independent methods – one based on classical density-functional theory and the other on the Smoluchowski equation – yield identical expressions for the equilibrium arm orientational distribution and numerical results suggest relatively weak orientational anisotropy. The resulting effective pair interactions between charged stars are fit with simple analytical functions.

¹Support from National Science Foundation grant DMR-0204020 is gratefully acknowledged.

Prefer Oral Session
 Prefer Poster Session

Hao Wang
hao.wang@ndsu.nodak.edu
Department of Physics, North Dakota State University

Date submitted: 23 Nov 2004

Electronic form version 1.4