

Abstract Submitted  
for the MAR05 Meeting of  
The American Physical Society

Sorting Category: 05.2 (E)

**Thermal Conductivity of  $\text{YBa}_2\text{Cu}_3\text{O}_7$**  DALE R. HARSHMAN, Physikon Research Corp., JOHN D. DOW, Arizona State University — It is widely, but incorrectly, believed that the thermal conductivity of  $\text{YBa}_2\text{Cu}_3\text{O}_7$  can only be explained in terms of a d-wave model. We show that an s-wave model of hole-pairing, combined with muon spectra that are observed to be s-wave, specific heat data with an observed zero-field linear-T term, superconductivity determined to be in the BaO layers, and non-superconducting  $\text{CuO}_2$ -plane bands describe the  $\text{YBa}_2\text{Cu}_3\text{O}_7$  data better than any d-wave model. Unlike any d-wave model, the s-wave explanation is also consistent with the superconductivity of Cu-doped  $\text{Sr}_2\text{YRuO}_6$  and  $\text{Ba}_2\text{YRuO}_6$ , at  $\approx 49$  K and  $\approx 83$  K, despite these compounds having no cuprate-planes, and the superconductivity of  $\text{GdSr}_2\text{Cu}_2\text{RuO}_8$  and  $\text{Gd}_{2-z}\text{Ce}_z\text{Sr}_2\text{Cu}_2\text{RuO}_{10}$  occurring at  $\approx 40$  K, despite the fact that their cuprate-planes are either antiferromagnetic or weakly ferromagnetic. In all of these compounds, the superconducting layer is the SrO or BaO layer. The errors which led to the widely believed erroneous concept of cuprate-plane superconductivity are discussed.

Prefer Oral Session  
 Prefer Poster Session

John D. Dow  
catsc@cox.net  
Arizona State University

Date submitted: 30 Nov 2004

Electronic form version 1.4