

Abstract Submitted
for the MAR05 Meeting of
The American Physical Society

Sorting Category: 05.4 (E)

Search for the Coexistence of Magnetism and Superconductivity in $\text{Ce}_{1-x}\text{Gd}_x\text{Ru}_2$ using the Mossbauer Effect¹

MICHAEL DE MARCO, RYAN HEARY, DERMOT COFFEY, Dept. of Physics, Buffalo State College, NY 14222, PE-CHUN HO, TODD SAYLES, BRIAN MAPLE, Dept. of Physics, University of California, San Diego, CA 92093, STEVE TOORONGIAN, MICHAEL HAKA, Nuclear Medicine Department, SUNY Buffalo, NY 14260 — We use the ⁹⁹Ru Mossbauer Effect (ME) to look for evidence for the coexistence of magnetism and superconductivity in $\text{Ce}_{1-x}\text{Gd}_x\text{Ru}_2$. These compounds have superconducting and ferromagnetic phases. The $T_c(x)$ and $\theta(x)$ curves cross at $x = 0.135$. Here $\theta(x)$ is the Curie temperature found by extrapolating magnetic susceptibility data below T_c . A sample of $\text{Ce}_{0.93}\text{Gd}_{0.07}\text{Ru}_2$ was prepared with enriched ⁹⁹Ru(97%) for which $T_c=4.5\text{K}$ and $\theta=4.19\text{K}$. The ME spectrum measured at 2.2K showed a pure quadrupole with a splitting of 0.36mm/s and an isomer shift of 0.12mm/s with no hyperfine field. The absence of the magnetic field in the sample brings into question the use of extrapolated magnetic susceptibility data as evidence for the coexistence of magnetism and superconductivity. Comparing the ME spectrum with that of CeRu_2 ($T_c=6.1\text{K}$) one sees that there is an impurity Ru phase. This suggests that previous evidence for magnetism below T_c on the basis of ⁵⁷Fe ME could be attributed to an impurity Fe phase.

¹Supported by the USDOE (DE-FG02-03ER46064)

Prefer Oral Session
 Prefer Poster Session

Michael De Marco
demarnj@buffalostate.edu

Date submitted: 08 Dec 2004

Electronic form version 1.4