

Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

Sorting Category: 13.6.4 (C)

Vibrations and thermal conductivity in inorganic and polymeric glasses. SERGEI SHENOGIN, Rensselaer Polytechnic Institute, ARUN BODAPATI, Rensselaer Polytechnic Institute, PAWEL KEBLINSKI, Rensselaer Polytechnic Institute — The mechanism of thermal transport in amorphous materials was studied by means of vibrational mode analysis and classical nonequilibrium molecular dynamics (MD) simulations. We studied four different model systems of (a) Lennard-Jones glass, (b) bead-spring model of an amorphous polymer, (c) amorphous silicon with Stillinger-Weber potential; and (d) all-atom model of glassy polystyrene with PCFF-type force field. For all structures we evaluated thermal conductivity from the harmonic theory of disordered solids [P.B.Allen, and J.L.Feldman, Phys.Rev.B 48, 12581 (1993)] and from direct MD simulations. We found that for all models but polystyrene, the harmonic theory accurately predicts thermal conductivity. By contrast, in the case of polystyrene, only $\sim 1/2$ of thermal conductivity can be explained within the harmonic approximation. Consequently, a major part of the transport has to be attributed to anharmonic coupling between vibrational modes. The reasons for the failure of harmonic theory of disordered solids to model amorphous glassy polymers will be discussed.

Prefer Oral Session
 Prefer Poster Session

Sergei Shenogin
shenos3@rpi.edu
Rensselaer Polytechnic Institute

Date submitted: 29 Nov 2005

Electronic form version 1.4