

Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

Sorting Category: 13.3 (C)

Composite-fermion antiparticle description of the hole excitation in the maximum-density droplet ALEV DEVRIM GUCLU, Cornell University, GUN SANG JEON, Pennsylvania State University, CYRUS J. UMRIGAR, Cornell University, JAINENDRA K. JAIN, Pennsylvania State University — The maximum-density droplet of quantum dots is a finite-size realization of the state at filling factor one. For a sufficiently small number of electrons, it becomes unstable to the creation of a central hole as the magnetic field is increased or the strength of the confinement potential reduced. The simplest model for the hole is a vortex at the center, which, however, is renormalized by edge excitations. We show that a remarkably accurate description of the actual hole state is achieved in terms of a “composite-fermion antiparticle,” which is surprising in view of the fact that composite fermions are thought to be relevant only in the fractional Hall regime. The composite-fermion antiparticle description also allows us to study the effect of Landau level mixing through variational and diffusion Monte Carlo calculations in a very efficient manner. Generalizations to systems containing several holes, as well as to the quasiholes of fractional quantum Hall states are presented.

Prefer Oral Session
 Prefer Poster Session

A. Devrim Guclu
guclud@tc.cornell.edu
Cornell University

Date submitted: 30 Nov 2005

Electronic form version 1.4